


# Introducing HyPeak: An international network on hydropeaking research, practice, and policy

Maria Alp<sup>1</sup>  | Ramon J. Batalla<sup>2,3</sup>  | Maria Dolores Bejarano<sup>4</sup>  |  
 Isabel Boavida<sup>5</sup>  | Hervé Capra<sup>1</sup>  | Mauro Carolli<sup>6</sup>  | Roser Casas-Mulet<sup>7,8</sup>  |  
 Maria João Costa<sup>5</sup>  | Jo Halvard Halleraker<sup>9,10</sup>  | Christoph Hauer<sup>11</sup>  |  
 Daniel S. Hayes<sup>12</sup>  | Atle Harby<sup>6</sup>  | Markus Noack<sup>13</sup>  | Antoni Palau<sup>2</sup>  |  
 Matthias Schneider<sup>14</sup>  | Lennart Schönfelder<sup>6</sup>  | Diego Tonolla<sup>15</sup>  |  
 Davide Vanzo<sup>16</sup>  | Terese Venus<sup>17</sup>  | Damià Vericat<sup>2,18</sup>  | Guido Zolezzi<sup>19</sup>  |  
 Maria Cristina Bruno<sup>20</sup> 

<sup>1</sup>RiverLy, INRAE, Villeurbanne, France

<sup>2</sup>Fluvial Dynamics Research Group, University of Lleida, Lleida, Spain

<sup>3</sup>Hydrological Processes Area, Catalan Institute for Water Research, Girona, Spain

<sup>4</sup>Natural Systems and Resources Department, Universidad Politécnica de Madrid, Madrid, Spain

<sup>5</sup>CERIS, Civil Engineering Research and Innovation for Sustainability, Instituto Superior Técnico, University of Lisbon, Lisbon, Portugal

<sup>6</sup>Energy Systems, SINTEF Energy Research, Trondheim, Norway

<sup>7</sup>Aquatic Systems Biology Unit, School of Life Sciences, Technical University of Munich, Freising, Germany

<sup>8</sup>Department of Infrastructure Engineering, The University of Melbourne, Melbourne, Victoria, Australia

<sup>9</sup>Department of Civil and Environmental Engineering, Norwegian University of Science and Technology, Trondheim, Norway

<sup>10</sup>Water Resource and Knowledge Department, Norwegian Environment Agency, Trondheim, Norway

<sup>11</sup>Institute of Hydraulic Engineering and River Research, University of Natural Resources and Life Sciences, Vienna, Austria

<sup>12</sup>Department of Water, Atmosphere and Environment, Institute of Hydrobiology and Aquatic Ecosystem Management, University of Natural Resources and Life Sciences, Vienna, Austria

<sup>13</sup>Institute of Applied Research, Karlsruhe University of Applied Science, Karlsruhe, Germany

<sup>14</sup>SJE Ecohydraulic Engineering GmbH, Stuttgart, Germany

<sup>15</sup>Institute of Natural Resource Sciences, Zurich University of Applied Sciences, Wädenswil, Switzerland

<sup>16</sup>Laboratory of Hydraulics, Hydrology and Glaciology, ETH Zürich, Zürich, Switzerland

<sup>17</sup>Technical University of Munich, Agricultural Production and Resource Economics Unit, Freising, Germany

<sup>18</sup>Landscape Dynamics and Biodiversity, Forest Sciences and Technology Centre of Catalonia, Solsona, Spain

<sup>19</sup>Department of Civil, Environmental and Mechanical Engineering, University of Trento, Trento, Italy

<sup>20</sup>Research and Innovation Centre, Fondazione Edmund Mach, San Michele all'Adige, Italy

## Correspondence

Maria Alp, INRAE, RiverLy, Villeurbanne, France.

Email: [maria.alp@inrae.fr](mailto:maria.alp@inrae.fr)

## Abstract

An increase in the demand for renewable energy is driving hydropower development and its integration with variable renewable energy sources. When hydropower is produced flexibly from hydropower plants, it causes rapid and frequent artificial flow fluctuations in rivers, a phenomenon known as hydropeaking. Hydropeaking and associated hydrological alterations cause multiple impacts on riverine habitats with cascading effects on ecosystem functioning and structure. Given the significance of

its ecological and socio-economic implications, mitigation of hydropeaking requires an inter- and transdisciplinary approach. An interdisciplinary network called HyPeak has been conceived to enrich international research initiatives and support hydropower planning and policy. HyPeak has been founded based on exchange and networking activities linking scientists from several countries where hydropeaking has been widespread for decades and numerous studies dedicated to the topic have been carried out. HyPeak aims to integrate members from other countries and continents in which hydropower production plays a relevant role, and grow to be a reference group that provides expert advice on the topic to policy-makers, as well as researchers, stakeholders, and practitioners in the field of hydropeaking.

#### KEYWORDS

anthropogenic impact, flow fluctuations, flow management, hydropower, interdisciplinary network, mitigation measures, rivers

## 1 | HYDROPEAKING AND HYDROPEAKED RIVERS

Hydropower is the largest source of renewable energy (71% as of 2016; Moran, Lopez, Moore, Müller, & Hyndman, 2018), generating 4,370 TWh with 1,330 GW of installed capacity globally in 2020 (IHA, 2021). Still, energy generation from hydropower must double by 2050 to meet the target of limiting global warming to 2°C (IEA, 2021). In addition to providing renewable energy, hydropower offers flexibility to the energy system at timescales ranging from seconds to seasons, which is important for the integration of variable renewable energy from wind and solar technologies (IEA, 2021). Depending on the site, technical layout, and operation mode, hydropower production may also have significant environmental and social impacts. When hydropower plants operate with rapid and frequent changes in energy generation to meet demands for flexibility, they cause rapid and frequent fluctuations of flow in outlets. Hydropeaking is the term used to describe rapid and frequent artificial flow fluctuations caused by flexible reservoir-operated hydropower production (Moog, 1993). The outlet flow may be delivered to reservoirs, lakes, or the sea, but the downstream effects are by far most pronounced when the outlets discharge directly to rivers, which is the focus of this paper.

Flow fluctuations can be characterised by changes in one or several flow regime components (i.e., flow magnitude, flow ratio, frequency, rate of change, and timing), which exceed those of the natural variability and intensity of the flow regime. Importantly, it is often the combination of these parameters that creates the most harmful effects (Harby & Noack, 2013; Hayes et al., 2019). Specifically, hydropeaking and associated hydrological alterations may result in thermal fluctuations (thermopeaking), geomorphological alterations (i.e., changes in sediment transport and in bed-material grain-size distribution), and changes in hydraulic habitat conditions (Table 1), the nature and extent of which will depend on the hydro-morphological characteristics of the downstream river (Vanzo et al., 2016). In turn, these changes cause numerous adverse impacts on aquatic biota and

ecosystem processes (Table 1). Furthermore, changes in hydrological conditions from hydropower may lead to socio-economic impacts related to water availability for irrigation, tourism, and recreational opportunities, the aesthetic value of rivers, and increased risks to local populations (Table 1).

As future demand for flexible hydropower operation is expected to increase, we must find ways of operating hydropower plants that limit negative impacts on downstream river ecosystems (Batalla et al., 2021; Jones, 2014). Currently available mitigation measures can be categorised into: (i) measures that directly modify the dynamics and the intensity of hydropeaking flow, (ii) indirect measures that reduce the impact on river habitat and hydro-morphological conditions, and (iii) complementary approaches (Hayes et al., 2022). The first group includes operational measures, such as lowering ramping rates by turbine flow restrictions, and structural measures, such as the construction of re-regulation reservoirs or compensation basins (Bruder et al., 2016). The second group of measures encompasses morphological improvements of the hydropeaked section through river restoration or addition of instream structures, as well as its reconnection with tributaries characterized by natural flow and sediment regime and/or presence of potential source populations (Baladrón, Costa, Bejarano, Pinheiro, & Boavida, 2021; Costa, Fuentes-Pérez, Boavida, Tuhtan, & Pinheiro, 2019; Greimel et al., 2018; Hauer et al., 2017). Finally, complementary measures include new approaches to coupling hydropeaking-based power production with alternative energy storage technologies (e.g., batteries, supercapacitors) or other renewables (i.e., solar, wind) (Haas, Nowak, & Palma-Behnke, 2019; Hayes et al., 2022).

A common understanding of sustainable hydropeaking mitigation has emerged in several European countries related to the implementation of the Water Framework Directive (Halleraker et al., 2016), but no common legal framework for hydropeaking thresholds exists. For instance, only the Alpine region has yet established legal regulations (i.e., on the national level in Austria and Switzerland and on the regional level in Bolzano, Italy). Some countries or regions (e.g., Spain

**TABLE 1** Main documented impacts of hydropeaking and list of representative recent papers for each impact (distributed according to different typologies)

| Type of impact | Impact   | Literature (examples)   |
|----------------|--|---|
| Physical       | Changes in temperature regime: Thermopeaking   | Carolli, Bruno, Siviglia, & Maiolini, 2012; Zolezzi, Siviglia, Toffolon, & Maiolini, 2011   |
|                | Riverbed clogging  | Hauer, Holzapfel, Tonolla, Habersack, & Zolezzi, 2019   |
|                | Changes in gas saturation: Saturopeaking   | Pulg, Wiik Vollset, Velle, & Stranzl, 2016  |
|                | Changes in underwater soundscapes: Soundpeaking  | Lumsdon et al., 2018  |
|                | Changes in grain-size distribution, bed-armouring and particles' mobility, and winnowing of sand and fine gravel | Béjar, Vericat, Batalla, & Gibbins, 2018; López, Garcia, Vericat, & Batalla, 2020; Vericat, Ville, Palau, & Batalla, 2020   |
|                | Changes in flow turbidity  | Béjar et al., 2018; Hauer et al., 2019; Lobera, Batalla, Vericat, López-Tarazón, & Tena, 2016   |
|                | Hydraulic impacts that is, increase in water velocity and turbulence, and sudden and rapid changes in flow depth | Hauer, Holzapfel, Leitner, & Graf, 2017; Shen & Diplas, 2010; Vanzo, Zolezzi, & Siviglia, 2016  |
| Biological     | Invertebrates: Drift, stranding, community composition, habitat use, egg mortality                               | Aksamit, Carolli, Vanzo, Weber, & Schmid, 2021; Bruno, Cashman, Maiolini, Biffi, & Zolezzi, 2016; Kennedy et al., 2016; Kjerstad, Arnekleiv, Speed, & Herland, 2018; Miller & Judson, 2014; Ruhí, Dong, McDaniel, Batzer, & Sabo, 2018; Schülting, Feld, Zeiringer, Hudek, & Graf, 2019 |
|                | Fish: Drift, stranding, egg mortality, population integrity, behaviour, habitat selection                        | Auer, Zeiringer, Führer, Tonolla, & Schmutz, 2017; Béjar et al., 2018; Boavida, Harby, Clarke, & Heggenes, 2017; Boavida et al., 2020; Capra et al., 2017; Casas-Mulet, Saltveit, & Alfredsen, 2015; Costa, Pinheiro, & Boavida, 2019; Hayes et al., 2019, 2021; Judes et al., 2020     |
|                | Periphyton: Reduced biomass, compositional changes, reduced nutritional quality                                  | Bondar-Kunze, Maier, Schönauer, Bahl, & Hein, 2016; Cashman, Harvey, Wharton, & Bruno, 2017   |
|                | Macrophytes and riparian vegetation: Communities composition, structure, and persistence                         | Bejarano, Jansson, & Nilsson, 2018; Bejarano, Sordo-Ward, Alonso, Jansson, & Nilsson, 2020; Gorla, Signarbieux, Turberg, Buttler, & Perona, 2015  |
|                | Foodwebs   | Holzapfel, Leitner, Habersack, Graf, & Hauer, 2017; Pearce et al., 2019   |
| Socio-economic | Risks to local populations   | Premstaller, Cavedon, Pisaturo, Schweizer, & Righetti, 2017; Venus et al., 2020   |
|                | Recreational opportunities (e.g., swimming, boating and fishing)   | Carolli et al., 2017; Venus et al., 2020; Venus & Sauer, 2022   |
|                | Aesthetic value of rivers  | Pflüger, Rackham, & Larned, 2010  |
|                | Water availability for irrigation  | Bryan et al., 2013  |

and the state of Baden-Württemberg in Germany) include hydropeaking mitigation recommendations in their river basin management plans, while others still work on a case-by-case basis (Moreira et al., 2019). With the exception of base flow requirements, which exist in many countries, the implementation of mitigation measures and thresholds for ramping restrictions are rare (Hayes et al., 2022; Tonolla, Bruder, & Schweizer, 2017).

## 2 | CHALLENGES AND NEEDS OF STUDYING HYDROPEAKED RIVERS

Given the complexity of transnational electricity markets and increasing electricity demand combined with the multifaceted impacts of hydropeaking, interdisciplinary, and transdisciplinary scientific research

is needed to support sustainable management and regulation of hydropeaking-based hydropower production. Developing such knowledge is challenging for several reasons: (i) the high and often unpredictable hydrological dynamism of mountain rivers, where hydropeaking typically occurs, renders field studies complex to carry out, and (ii) the highly variable geographic and geomorphological settings, technical approaches, and legal regulations adopted at national and regional scales limit the transferability of results. Finally, (iii) stronger links between disciplines are urgently needed, but have yet been poorly developed. This concerns in particular, the social sciences, which should be integrated into biophysical research on hydropeaking (e.g., risks to local populations, public acceptance, power markets projections based on climate change). At the same time, an increasing number of research tools and decision support tools have been developed in recent years and offer new approaches to key disciplines

**TABLE 2** Novel research tools expanding the opportunities in the research on hydropeaking

| Aspects assessed                      | Tools  | Some representative references (illustrating application in the field of hydropeaking)  |
|---------------------------------------|--|---|
| Physical processes                    | Hydraulic, hydrological and topographical modelling including simulations, particle tracking techniques, assessment of river topography and morphology through geomatics, thermal imagery, remote sensing, experimentation in hydraulic laboratories | Brasington, Vericat, & Rychkov, 2012; Carolli et al., 2015; Sauterleute & Charmasson, 2014  |
| Biology                               | Habitat and population modelling; underwater cameras and telemetry for behavioral observations; experimentation in hydraulic laboratories and open-air flumes; impacts classification systems  | Bakken et al., 2021; Boavida et al., 2021; Bruno et al., 2016; Capra et al., 2017; Costa et al., 2019b; Hedger et al., 2018; Holzapfel et al., 2017; Leitner, Hauer, & Graf, 2017; Person, Bieri, Peter, & Schleiss, 2014; Schneider et al., 2017; Schülting et al., 2018 |
| Optimising hydropower plant operation | Market models, power system analyses, decision support tools   | Kong, Skjelbred, & Fosso, 2020  |

related to hydropeaking and to hydropower production planning (Table 2).

In this context, several tasks appear to be of high research priority for the scientific community working on the topic of hydropeaking:

1. Compile an overview of the localization and typology of hydropeaking at a large scale (continental). For instance, no geo-database including source and severity of hydropeaking impacts across Europe exists. Standardization of tools used for characterizing the hydropeaking hydrological regime would be necessary for this task.
2. Identify the most informative indicators for assessing hydropeaking impacts on rivers. While a number of biophysical indicators have already been suggested for this purpose (e.g., water temperature, sediment transport, invertebrate drift, fish stranding; see Bruder et al., 2016), further indicators, covering other types of organisms and life stages, levels of biological organization (e.g., food webs), ecosystem functions and associated services (e.g., carbon retention and greenhouse gas emissions), geomorphological processes (e.g., river-bed degradation, bed-material winnowing, armouring, habitat degradation), and societal needs (e.g., recreational opportunities), need to be developed.
3. Continue development of technical approaches to limit the negative impacts of hydropower plant operation on river ecosystems and of procedures allowing to select the most locally-appropriate ones, evaluate their feasibility by implementing cost-benefit (ecological, social, etc.) analyses and assess their efficiency.
4. Elaborate inter- and transdisciplinary approaches for finding compromises between hydropower plant operation and ecological sustainability, to achieve an economic sustainability of mitigation measures and carbon-free energy production, in view of the pressures of energy markets, private and public interests, as well as the challenges imposed by climate change. This issue needs to be addressed in a general framework of energy planning and requires increasing the flexibility in the power system and coupling different renewable energy sources.


5. Create and reinforce functioning links between research and application (researchers of various disciplines, legislators, public river managers, hydropower producers). This implies an efficient reciprocal transfer of knowledge and tools between researchers and practitioners to support the elaboration of informed policies, the choice of technical solutions, and the assessment of their efficiency at different scales (local, river basin, national, international).

Several of the listed tasks require a systematic inter- and transdisciplinary approach. Moreover, exchanging tools and experiences across countries appears crucial to advance towards sustainable hydropeaking management on a large scale (e.g., Pittman, Tiessen, & Montaña, 2016). International and interdisciplinary networks are essential for improving the transfer of knowledge across borders and from science to practice. In Europe, several recent networks and projects (e.g., smires.eu, converges.eu, FitHydro.eu) have succeeded in creating key scientific consortia in aquatic environmental research, building strong links across borders and disciplines, and supporting the development of environmental policies (e.g., Brils, 2020). In the case of hydropeaking, hydropower operators, public decision-makers and stakeholders must be involved in such networks to promote knowledge-sharing and dialogue concerning the degree of hydropeaking impacts on the environment and the selection of the best cost-effective mitigation measures (Barillier, Beche, Malavoi, & Gouraud, 2021).

### 3 | FOUNDING AND ACTIVITIES OF THE HYPEAK NETWORK

The interdisciplinary network on Hydropeaking Research (HyPeak) was founded after an interactive and well-attended webinar held during the 13th International Symposium on Ecohydraulics in November 2020, to propose a framework for crossing interdisciplinary perspectives on hydropeaking (Figure 1). In 2021, the network started to

**FIGURE 1** A schematic representation of the multidisciplinary approach required to tackle hydropeaking impacts on the different components of socio-ecosystems and propose mitigation strategies [Color figure can be viewed at [wileyonlinelibrary.com](http://wileyonlinelibrary.com)]



operate with the core group of European researchers covering a wide range of disciplines (hydrology, geomorphology, hydraulic engineering, ecology, economics) and countries. The first steps consisted in reaching an agreement on the exact mission of the network, its organization, and enrollment/operational rules indispensable before opening it up to interested future members. This resulted in a formalised HyPeak Charter (<https://www.researchgate.net/project/HyPeak-Hydropeaking-Research-Network>), which identified the overall goal of HyPeak as one of becoming a reference group of experts to support evidence-based guidelines and legislation at different levels (national to international) and enhance the value of the interaction between research and policy. Therefore, our activities focus on collaborations among researchers and the creation of strong bridges with practitioners (hydropower producers, environmental agencies, consultants, or legislators). Our key mission is to stimulate integrative hydropeaking research across disciplines to support:

1. Assessment of environmental effects and related socio-economic issues of hydropeaking at various spatial and temporal scales;
2. Fundamental understanding of hydropeaking dynamics and related biophysical processes;
3. Improvements of mitigation measures and management strategies;

4. Promotion of environmentally sustainable approaches to hydropeaking;
5. Providing recommendations for national and international policies and supporting their integration.

Involvement in the HyPeak Network is possible at several levels: (i) as a General Member: adhering to the principles of the network and following its activities via email newsletter; (ii) as a Core Group Member: getting actively involved in specific activities of the network (see below); (iii) as a member of the Executive Board: centralizing the activities of the network and assuring continuity of the Core Group meetings. Importantly, our choice is to keep the HyPeak network independent of any funding other than funding for scientific research.

We envisage a wide range of activities according to the above-stated key mission in the upcoming years aiming at:

- (i) Keeping the members informed of news in relation to hydropeaking across countries and research fields (new legislation, new research results, conferences, etc.);
- (ii) Preparing collaborative international research initiatives to optimise acquisition and dissemination of knowledge (co-supervision of doctoral students, conducting parallel experimental approaches

across continents/hydro-bioregions, working on transboundary study cases, sharing/enlarging datasets for cross-continental or global impact analyses);

- (iii) Exchanging knowledge and enhancing the common understanding across disciplines (e.g., river ecology and economics);
- (iv) Fostering the involvement of managers in defining research guidelines;
- (v) Expanding the geographic coverage of the network well outside of Europe.

Analysing current knowledge gaps and key expectations of practitioners and legislators towards the scientific community belongs to our key short-term objectives. Therefore, one of HyPeak's first activities consisted in conducting an extensive online survey targeting hydropower stakeholders and researchers working on the topic of hydropeaking (released in December 2021). Together with a follow-up Delphi study, the aims of this work are to gather global research priorities, emerging issues related to hydropeaking impacts and their mitigation, and to analyse stakeholder perceptions of the hydropeaking effects on the ecosystem services in rivers. Key research questions relevant for the practitioners identified through this survey will allow us to adjust the baseline for future activities of the HyPeak network and beyond, such as the prioritization of future research to meet the policy and management needs of hydropeaked rivers. In addition, the lessons from the stakeholder perceptions of the ecosystem service effects of hydropeaking will support science, governance, and management in the effective communication of impacts and necessary mitigation approaches.

The preparation of the special issue "Innovations in Hydropeaking Research" in *River Research & Applications*, led by an editorial board of HyPeak Core Group members represents another activity fostering communication of current research in the field. In the same line, a series of open webinars on sustainable hydropeaking is organised by HyPeak starting from 2022. The idea is to present both researcher and practitioner perspectives on hydropeaking through a selection of speakers with different professional backgrounds and geographical origins.

## 4 | CONCLUSIONS

Our current knowledge on how many rivers are exposed to hydropeaking across continents, the severity of the resulting impacts, and their dependence on the specific boundary conditions (river type, geographic setting, management scheme, or history of hydropeaking), is currently extremely limited. A joint effort enabled by an international network in the long term could help collect, standardise, and share data on hydropeaked rivers (hydrological, geomorphological, ecological, legal, social...) on a large scale. These data could be used, for instance, to produce maps of hydropeaked rivers across continents and allow us to have a global overview of the hydropeaking phenomenon and status updates about ecological effective mitigation measures. Similarly, a future task would be to launch comparisons and knowledge transfer across borders in terms of hydropeaking extent

and impact, the weight of this type of energy production in the energy mix, legislation, monitoring tools, and mitigation approaches as well as public perception.

As stated in the HyPeak Charter, our network is independent and based on the voluntary contributions of its members. The HyPeak Network is open to new members interested in participating in it, at the desired level of commitment as stated in Annex 1 of the Charter. The first step for any interested person is to enrol as a General Member by filling the form available at <https://forms.gle/CsahiE6dWB5SuLaAA>.

## ACKNOWLEDGEMENTS

The authors thank the organisers of the 13th International Symposium on Ecohydraulics (ISE2020) for the opportunity of presenting the HyPeak Network in an online workshop on November 23, 2020.

## DATA AVAILABILITY STATEMENT

Data sharing is not applicable to this article as no new data were created or analyzed in this study.

## ORCID

Maria Alp  <https://orcid.org/0000-0003-4295-6361>

Ramon J. Batalla  <https://orcid.org/0000-0001-8454-2314>

María Dolores Bejarano  <https://orcid.org/0000-0002-5180-4442>

Isabel Boavida  <https://orcid.org/0000-0001-6005-3751>

Hervé Capra  <https://orcid.org/0000-0002-3965-4314>

Mauro Carolli  <https://orcid.org/0000-0001-9625-8441>

Roser Casas-Mulet  <https://orcid.org/0000-0002-7139-8859>

Maria João Costa  <https://orcid.org/0000-0001-6395-6868>

Jo Halvard Halleraker  <https://orcid.org/0000-0002-6031-7726>

Christoph Hauer  <https://orcid.org/0000-0001-8704-2198>

Daniel S. Hayes  <https://orcid.org/0000-0003-3709-8630>

Atle Harby  <https://orcid.org/0000-0002-4965-6970>

Markus Noack  <https://orcid.org/0000-0002-3625-0357>

Antoni Palau  <https://orcid.org/0000-0001-7346-3936>

Matthias Schneider  <https://orcid.org/0000-0002-8322-6974>

Lennart Schönfelder  <https://orcid.org/0000-0002-0151-2812>

Diego Tonolla  <https://orcid.org/0000-0002-1172-0033>

Davide Vanzo  <https://orcid.org/0000-0002-2033-9197>

Terese Venus  <https://orcid.org/0000-0003-0729-2416>

Damià Vericat  <https://orcid.org/0000-0002-5685-4895>

Guido Zolezzi  <https://orcid.org/0000-0003-2807-7387>

Maria Cristina Bruno  <https://orcid.org/0000-0001-7860-841X>

## REFERENCES

- Aksamit, C. K., Carolli, M., Vanzo, D., Weber, C., & Schmid, M. (2021). Macroinvertebrate recovery to varying hydropeaking frequency: A small hydropower plant experiment. *Frontiers in Environmental Science*, 8, 602374. <https://doi.org/10.3389/fenvs.2020.602374>
- Auer, S., Zeiringer, B., Führer, S., Tonolla, D., & Schmutz, S. (2017). Effects of river bank heterogeneity and time of day on drift and stranding of juvenile European grayling (*Thymallus thymallus* L.) caused by hydropeaking. *Science of the Total Environment*, 575, 1515–1521. <https://doi.org/10.1016/j.scitotenv.2016.10.029>



- Bakken, T. H., Harby, A., Forseth, T., Ugedal, O., Sauterleute, J. F., Halleraker, J. H., & Alfredsen, K. (2021). Classification of hydropeaking impacts on Atlantic salmon populations in regulated rivers. *River Research and Applications*, 1–13. <https://doi.org/10.1002/rra.3917>
- Baladrón, A., Costa, M. J., Bejarano, M. D., Pinheiro, A., & Boavida, I. (2021). Can vegetation provide shelter to cyprinid species under hydropeaking? *Science of the Total Environment*, 769, 145339. <https://doi.org/10.1016/j.scitotenv.2021.145339>
- Barillier, A., Beche, L., Malavoi, J.-R., & Gouraud, V. (2021). Identification of effective hydropeaking mitigation measures: Are hydraulic habitat models sufficient in a global approach? *Journal of Ecohydraulics*, 6, 172–185. <https://doi.org/10.1080/24705357.2020.1856008>
- Batalla, R. J., Gibbins, C. N., Alcázar, J., Brasington, J., Buendia, C., Garcia, C., ... Vericat, D. (2021). Hydropeaked rivers need attention. *Environmental Research Letters*, 16, 02100. <https://doi.org/10.1088/1748-9326/abce26>
- Béjar, M., Vericat, D., Batalla, R. J., & Gibbins, C. N. (2018). Variation in flow and suspended sediment transport in a montane river affected by hydropeaking and instream mining. *Geomorphology*, 310, 69–83. <https://doi.org/10.1016/j.geomorph.2018.03.001>
- Bejarano, M. D., Jansson, R., & Nilsson, C. (2018). The effects of hydropeaking on riverine plants: A review. *Biological Reviews of the Cambridge Philosophical Society*, 93, 658–673. <https://doi.org/10.1111/brv.12362>
- Bejarano, M. D., Sordo-Ward, Á., Alonso, C., Jansson, R., & Nilsson, C. (2020). Hydropeaking affects germination and establishment of river-bank vegetation. *Ecological Applications*, 30, e02076. <https://doi.org/10.1002/eap.2076>
- Boavida, I., Ambrósio, F., Costa, M. J., Quesada, A., Portela, M. M., Pinheiro, A., & Godinho, F. (2020). Habitat use by *Pseudochondrostoma duriense* and *Squalius carolitertii* downstream of a small-scale hydro-power plant. *Water*, 12, 2522. <https://doi.org/10.3390/w12092522>
- Boavida, I., Costa, M. J., Portela, M. M., Godinho, F., Tuhtan, J., & Pinheiro, A. (2021). Do cyprinid fish use lateral flow-refuges during hydropeaking? *River Research and Applications*, 1–7. <https://doi.org/10.1002/rra.3863>
- Boavida, I., Harby, A., Clarke, K. D., & Heggenes, J. (2017). Move or stay: Habitat use and movements by Atlantic salmon parr (*Salmo salar*) during induced rapid flow variations. *Hydrobiologia*, 785, 261–275. <https://doi.org/10.1007/s10750-016-2931-3>
- Bondar-Kunze, E., Maier, S., Schönauer, D., Bahl, N., & Hein, T. (2016). Antagonistic and synergistic effects on a stream periphyton community under the influence of pulsed flow velocity increase and nutrient enrichment. *Science of the Total Environment*, 573, 594–602. <https://doi.org/10.1016/j.scitotenv.2016.08.158>
- Brasington, J., Vericat, D., & Rychkov, I. (2012). Modeling river bed morphology, roughness, and surface sedimentology using high resolution terrestrial laser scanning. *Water Resources and Research*, 48, W11519. <https://doi.org/10.1029/2012WR012223>
- Brils, J. (2020). Including sediment in European River basin management plans: Twenty years of work by SedNet. *Journal of Soils and Sediments*, 20, 4229–4237. <https://doi.org/10.1007/s11368-020-02782-1>
- Bruder, A., Tonolla, D., Schweizer, S. P., Vollenweider, S., Langhans, S. D., & Wüest, A. (2016). A conceptual framework for hydropeaking mitigation. *Science of the Total Environment*, 568, 1204–1212. <https://doi.org/10.1016/j.scitotenv.2016.05.032>
- Bruno, M. C., Cashman, M. J., Maiolini, B., Biffi, S., & Zolezzi, G. (2016). Responses of benthic invertebrates to repeated hydropeaking in semi-natural flume simulations. *Ecohydrology*, 9, 68–82. <https://doi.org/10.1002/eco.1611>
- Bryan, B. A., Higgins, A., Overton, I. C., Holland, K., Lester, R. E., King, D., ... Kirby, M. (2013). Ecohydrological and socioeconomic integration for the operational management of environmental flows. *Ecological Applications*, 23, 999–1016. <https://doi.org/10.1890/12-2104.1>
- Capra, H., Plichard, L., Bergé, J., Pella, H., Ovidio, M., McNeil, E., & Lamouroux, N. (2017). Fish habitat selection in a large hydropeaking river: Strong individual and temporal variations revealed by telemetry. *Science of the Total Environment*, 578, 109–120. <https://doi.org/10.1016/j.scitotenv.2016.10.155>
- Carolli, M., Bruno, M. C., Siviglia, A., & Maiolini, B. (2012). Responses of benthic invertebrates to abrupt changes of temperature in flume simulations. *River Research and Applications*, 28, 678–691. <https://doi.org/10.1002/rra.1520>
- Carolli, M., Vanzo, D., Siviglia, A., Zolezzi, G., Bruno, M. C., & Alfredsen, K. (2015). A simple procedure for the assessment of hydropeaking flow alterations applied to several European streams. *Aquatic Sciences*, 77, 639–653. <https://doi.org/10.1007/s00027-015-0408-5>
- Carolli, M., Zolezzi, G., Geneletti, D., Siviglia, A., Carolli, F., & Cainelli, O. (2017). Modelling white-water rafting suitability in a hydropower regulated Alpine River. *Science of the Total Environment*, 579, 1035–1059. <https://doi.org/10.1016/j.scitotenv.2016.11.049>
- Casas-Mulet, R., Saltveit, S. J., & Alfredsen, K. (2015). The survival of Atlantic salmon (*Salmo salar*) eggs during dewatering in a river subjected to hydropeaking. *River Research and Applications*, 31, 433–446. <https://doi.org/10.1002/rra.2827>
- Cashman, M. J., Harvey, G. L., Wharton, G., & Bruno, M. C. (2017). Wood mitigates the effect of hydropeaking scour on periphyton biomass and nutritional quality in semi-natural flume simulations. *Aquatic Sciences*, 79, 459–471. <https://doi.org/10.1007/s00027-016-0510-3>
- Costa, M. J., Fuentes-Pérez, J. F., Boavida, I., Tuhtan, J. A., & Pinheiro, A. N. (2019a). Fish under pressure: Examining behavioural responses of Iberian barbel under simulated hydropeaking with instream structures. *PLoS One*, 14, e0211115. <https://doi.org/10.1371/journal.pone.0211115>
- Costa, M. J., Pinheiro, A., & Boavida, I. (2019b). Habitat enhancement solutions for Iberian cyprinids affected by hydropeaking: Insights from flume research. *Sustainability*, 11, 6998. <https://doi.org/10.3390/su11246998>
- Gorla, L., Signarbieux, C., Turberg, P., Buttler, A., & Perona, P. (2015). Effects of hydropeaking waves' offsets on growth performances of juvenile *Salix* species. *Ecological Engineering*, 77, 297–306. <https://doi.org/10.1016/j.ecoleng.2015.01.019>
- Greimel, F., Schülting, L., Graf, W., Bondar-Kunze, E., Auer, S., Zeiringer, B., & Hauer, C. (2018). Hydropeaking impacts and mitigation. In S. Schmutz & J. Sendzimir (Eds.), *Riverine ecosystem management. Aquatic ecology series* (Vol. 8, pp. 91–110). Springer, Cham, Switzerland. [https://doi.org/10.1007/978-3-319-73250-3\\_5](https://doi.org/10.1007/978-3-319-73250-3_5)
- Haas, J., Nowak, W., & Palma-Behnke, R. (2019). Multi-objective planning of energy storage technologies for a fully renewable system: Implications for the main stakeholders in Chile. *Energy Policy*, 126, 494–506. <https://doi.org/10.1016/j.enpol.2018.11.034>
- Halleraker, J. H., van de Bund, W., Bussetini, M., Gosling, R., Döbbelt-Grüne, S., Hensman, J., ... Döbbelt-Grüne, S. (2016). Working group ECOSTAT report on common understanding of using mitigation measures for reaching good ecological potential for heavily modified water bodies - part 1: Impacted by water storage. *JRC Report EUR*, 28413, 1–103. <https://doi.org/10.2760/649695>
- Harby, A., & Noack, M. (2013). Rapid flow fluctuations and impacts on fish and the aquatic ecosystem. In I. Maddock, A. Harby, P. Kemp, & P. Wood (Eds.), *Ecohydraulics: An integrated approach* (pp. 323–335). Oxford: John Wiley & Sons.
- Hauer, C., Holzapfel, P., Leitner, P., & Graf, W. (2017). Longitudinal assessment of hydropeaking impacts on various scales for an improved process understanding and the design of mitigation measures. *Science of the Total Environment*, 575, 1503–1514. <https://doi.org/10.1016/j.scitotenv.2016.10.031>
- Hauer, C., Holzapfel, P., Tonolla, D., Habersack, H., & Zolezzi, G. (2019). In situ measurements of fine sediment infiltration (FSI) in gravel-bed

- rivers with a hydropeaking flow regime. *Earth Surface Processes and Landforms*, 44, 433–448. <https://doi.org/10.1002/esp.4505>
- Hayes, D. S., Lautsch, E., Unfer, G., Greimel, F., Zeiringer, B., Höller, N., & Schmutz, S. (2021). Response of European grayling, *Thymallus thymallus*, to multiple stressors in hydropeaking rivers. *Journal of Environmental Management*, 292, 112737. <https://doi.org/10.1016/j.jenvman.2021.112737>
- Hayes, D. S., Moreira, M., Boavida, I., Haslauer, M., Unfer, G., Zeiringer, B., Greimel, F., Auer, S., Ferreira, T., & Schmutz, S. (2019). Life stage-specific hydropeaking flow rules. *Sustainability*, 11(6), 1547. <https://doi.org/10.3390/su11061547>
- Hayes, D. S., Schülting, L., Carolli, M., Greimel, F., Batalla, R. J., & Casas-Mulet, R. (2022). Hydropeaking: Processes, effects, and mitigation. *Reference Module in Earth Systems and Environmental Sciences*, Elsevier (pp. 134–149). <https://doi.org/10.1016/B978-0-12-819166-8.00171-7>
- Hedger, R. D., Sauterleute, J., Sundt Hansen, L. E., Forseth, T., Ugedal, O., Diserud, O. H., & Bakken, T. H. (2018). Modelling the effect of hydropeaking induced stranding mortality on Atlantic salmon population abundance. *Ecohydrology*, 11(5), e1960. <https://doi.org/10.1002/eco.1960>
- Holzapfel, P., Leitner, P., Habersack, H., Graf, W., & Hauer, C. (2017). Evaluation of hydropeaking impacts on the food web in alpine streams based on modelling of fish- and macroinvertebrate habitats. *Science of the Total Environment*, 575, 1489–1502. <https://doi.org/10.1016/j.scitotenv.2016.10.016>
- International Energy Agency (IEA). (2021). *Hydropower Special Market Report. Analysis and forecast to 2030*. <https://www.iea.org/reports/hydropower-special-market-report>
- International Hydropower Association (IHA). (2021). *Hydropower status report 2021*. London, UK: International Hydropower Association Limited. <https://www.hydropower.org/publications/2021-hydropower-status-report>
- Jones, N. E. (2014). The dual nature of hydropeaking rivers: Is ecopeaking possible? *River Research and Applications*, 30, 521–526. <https://doi.org/10.1002/rra.2653>
- Judes, C., Gouraud, V., Capra, H., Maire, A., Barillier, A., & Lamouroux, N. (2020). Consistent but secondary influence of hydropeaking on stream fish assemblages in space and time. *Journal of Ecohydraulics*, 6, 157–171. <https://doi.org/10.1080/24705357.2020.1790047>
- Kennedy, T. A., Muehlbauer, J. D., Yackulic, C. B., Lytle, D. A., Miller, S. W., Dibble, K. L., ... Baxter, C. V. (2016). Flow management for hydropeaking extirpates aquatic insects, undermining river food webs. *Bioscience*, 66, 561–575. <https://doi.org/10.1093/biosci/biw059>
- Kjærstad, G., Arnekleiv, J. V., Speed, J. D. M., & Herland, A. K. (2018). Effects of hydropeaking on benthic invertebrate community composition in two central Norwegian rivers. *River Research and Applications*, 34, 218–231. <https://doi.org/10.1002/rra.3241>
- Kong, J., Skjelbred, H. I., & Fosso, O. B. (2020). An overview on formulations and optimization methods for the unit-based short-term hydro scheduling problem. *Electric Power Systems Research*, 178, 106027. <https://doi.org/10.1016/j.epsr.2019.106027>
- Leitner, P., Hauer, C., & Graf, W. (2017). Habitat use and tolerance levels of macroinvertebrates concerning hydraulic stress in hydropeaking rivers—A case study at the Ziller River in Austria. *Science of the Total Environment*, 575, 112–118. <https://doi.org/10.1016/j.scitotenv.2016.10.011>
- Lobera, G., Batalla, R. J., Vericat, D., López-Tarazón, J. A., & Tena, A. (2016). Sediment transport in two Mediterranean regulated rivers. *Science of the Total Environment*, 540, 101–113. <https://doi.org/10.1016/j.scitotenv.2015.08.018>
- López, R., Garcia, C., Vericat, D., & Batalla, R. J. (2020). Downstream changes of particle entrainment in a hydropeaked river. *Science of the Total Environment*, 745, 140952. <https://doi.org/10.1016/j.scitotenv.2020.140952>
- Lumsdon, I. A., Bruno, M. C., Righetti, M., Tockner, K., Tonolla, D., & Zarfl, C. (2018). Soundpeaking – Hydropeaking induced changes in river soundscapes. *River Research and Applications*, 34(1), 3–12. <https://doi.org/10.1002/rra.3229>
- Miller, S. W., & Judson, S. (2014). Responses of macroinvertebrate drift, benthic assemblages, and trout foraging to hydropeaking. *Canadian Journal of Fisheries and Aquatic Sciences*, 687, 675–687. <https://doi.org/10.1139/cjfas-2013-0562>
- Moog, O. (1993). Quantification of daily peak hydropower effects on aquatic fauna and management to minimize environmental impacts. *Regulated Rivers: Research & Management*, 8, 5–14. <https://doi.org/10.1002/rrr.3450080105>
- Moran, E. F., Lopez, M. C., Moore, N., Müller, N., & Hyndman, D. W. (2018). Sustainable hydropower in the 21<sup>st</sup> century. *Proceedings of the National Academy of Sciences*, 115, 11891–11898. <https://doi.org/10.1073/pnas.1809426115>
- Moreira, M., Hayes, D. S., Boavida, I., Schletterer, M., Schmutz, S., & Pinheiro, A. (2019). Ecologically-based criteria for hydropeaking mitigation: A review. *Science of the Total Environment*, 657, 1508–1522. <https://doi.org/10.1016/j.scitotenv.2018.12.107>
- Pearce, J. L., Smokorowski, K. E., Brush, J., Timusk, E., Marty, J., & Power, M. (2019). Unrestricted ramping rates and long-term trends in the food web metrics of a boreal river. *River Research and Applications*, 35, 1575–1589. <https://doi.org/10.1002/rra.3516>
- Person, E., Bieri, M., Peter, A., & Schleiss, A. J. (2014). Mitigation measures for fish habitat improvement in Alpine rivers affected by hydropower operations. *Ecohydrology*, 7(2), 580–599. <https://doi.org/10.1002/eco.1380>
- Pflüger, Y., Rackham, A., & Larned, S. (2010). The aesthetic value of river flows: An assessment of flow preferences for large and small rivers. *Landscape and Urban Planning*, 95, 68–78. <https://doi.org/10.1016/j.landurbplan.2009.12.004>
- Pittman, J., Tiessen, H., & Montaña, E. (2016). The evolution of interdisciplinarity over 20 years of global change research by the IAI. *Current Opinion in Environmental Sustainability*, 19, 87–93. <https://doi.org/10.1016/j.cosust.2015.12.004>
- Premstaller, G., Cavedon, V., Pisaturo, G. R., Schweizer, S., & Righetti, M. (2017). Hydropeaking mitigation project on a multi-purpose hydro-scheme on Valsura River in South Tyrol/Italy. *Science of the Total Environment*, 574, 642–653. <https://doi.org/10.1016/j.scitotenv.2016.09.088>
- Pulg, U., Wiik Vollset, K., Velle, G., & Stranzl, S. (2016). First observations of saturopeaking: Characteristics and implications. *Science of the Total Environment*, 573, 1615–1621. <https://doi.org/10.1016/j.scitotenv.2016.09.143>
- Ruhi, A., Dong, X., McDaniel, C. H., Batzer, D. P., & Sabo, J. L. (2018). Detrimental effects of a novel flow regime on the functional trajectory of an aquatic invertebrate metacommunity. *Global Change Biology*, 24, 3749–3765. <https://doi.org/10.1111/gcb.14133>
- Sauterleute, J. F., & Charmasson, J. (2014). A computational tool for the characterisation of rapid fluctuations in flow and stage in rivers caused by hydropeaking. *Environmental Modelling & Software*, 55, 266–278. <https://doi.org/10.1016/j.envsoft.2014.02.004>
- Schneider, M., Kopecki, I., Tuhtan, J., Sauterleute, J. F., Zinke, P., Bakken, T. H., ... Merigoux, S. (2017). A fuzzy rule based model for the assessment of macrobenthic habitats under hydropeaking impact. *River Research and Applications*, 33(3), 377–387.
- Schülting, L., Feld, C. K., Zeiringer, B., Hudek, H., & Graf, W. (2019). Macroinvertebrate drift response to hydropeaking: An experimental approach to assess the effect of varying ramping velocities. *Ecohydrology*, 12, e2032. <https://doi.org/10.1002/eco.2032>
- Shen, Y., & Diplas, P. (2010). Modeling unsteady flow characteristics of hydropeaking operations and their implications on fish habitat. *Journal of Hydraulic Engineering*, 136, 1053–1066. [https://doi.org/10.1061/\(ASCE\)HY.1943-7900.0000112](https://doi.org/10.1061/(ASCE)HY.1943-7900.0000112)



- Tonolla, D., Bruder, A., & Schweizer, S. (2017). Evaluation of mitigation measures to reduce hydropeaking impacts on river ecosystems – A case study from the Swiss Alps. *Science of the Total Environment*, 574, 594–604. <https://doi.org/10.1016/j.scitotenv.2016.09.101>
- Vanzo, D., Zolezzi, G., & Siviglia, A. (2016). Eco-hydraulic modelling of the interactions between hydropeaking and river morphology. *Ecohydrology*, 9, 421–437. <https://doi.org/10.1002/eco.1647>
- Venus, T. E., Hinzmann, M., Bakken, T. H., Gerdes, H., Godinho, F. N., Hansen, B., Pinheiro A., & Sauer, J. (2020). The public's perception of run-of-the-river hydropower across Europe. *Energy Policy*, 140, 111422. <https://doi.org/10.1016/j.enpol.2020.111422>
- Venus, T. E., & Sauer, J. (2022). Certainty pays off: The public's value of environmental monitoring. *Ecological Economics*, 191, 107220. <https://doi.org/10.1016/j.ecolecon.2021.107220>
- Vericat, D., Ville, F., Palau, A., & Batalla, R. J. (2020). Effects of hydropeaking on bed mobility: Evidence from a Pyrenean river. *Water*, 12, 178. <https://doi.org/10.3390/w12010178>
- Zolezzi, G., Siviglia, A., Toffolon, M., & Maiolini, B. (2011). Thermo peaking in alpine streams: Event characterization and time scales. *Ecohydrology*, 4, 564–576. <https://doi.org/10.1002/eco.132>

**How to cite this article:** Alp, M., Batalla, R. J., Bejarano, M. D., Boavida, I., Capra, H., Carolli, M., Casas-Mulet, R., Costa, M. J., Halleraker, J. H., Hauer, C., Hayes, D. S., Harby, A., Noack, M., Palau, A., Schneider, M., Schönfelder, L., Tonolla, D., Vanzo, D., Venus, T., ... Bruno, M. C. (2022). Introducing HyPeak: An international network on hydropeaking research, practice, and policy. *River Research and Applications*, 1–9. <https://doi.org/10.1002/rra.3996>